

This is a repository copy of *Viral host-adaptation : insights from evolution experiments with phages*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/77996/>

Version: Submitted Version

Article:

Hall, James Pj, Harrison, Ellie and Brockhurst, Michael A orcid.org/0000-0003-0362-820X
(2013) *Viral host-adaptation : insights from evolution experiments with phages*. *Current Opinion in Virology*. pp. 572-577. ISSN 1879-6265

<https://doi.org/10.1016/j.coviro.2013.07.001>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Viral host-adaptation: insights from evolution experiments with phages

James P. J. Hall, Ellie Harrison, Michael A. Brockhurst

Department of Biology, University of York, Wentworth Way, York YO10 5DD

Author for correspondence:

Professor Michael A. Brockhurst

Postal address: Department of Biology, University of York, Wentworth Way, York YO10 5DD

Telephone: 00 44 1904 328 576

Email: michael.brockhurst@york.ac.uk

Abstract

Phages, viral parasites of bacteria, share fundamental features of pathogenic animal and plant viruses and represent a highly tractable empirical model system to understand viral evolution and in particular viral host-adaptation. Phage adaptation to a particular host genotype often results in improved fitness by way of parallel evolution whereby independent lineages hit upon identical adaptive solutions. By contrast, phage adaptation to an evolving host population leads to the evolution of increasing host-range over time and correlated phenotypic and genetic divergence between populations. Phage host-range expansion frequently occurs by a process of stepwise evolution of multiple mutations, and host-shifts are often constrained by mutational availability, pleiotropic costs or ecological conditions.

Published in Current Opinion in Virology 3:572-7 doi: 10.1016/j.coviro.2013.07.001

Introduction

Evolutionary studies of phages, viral parasites of bacteria, can reveal general principles of viral biology because phages share many of the fundamental features of pathogenic animal and plant viruses [1]. Large populations of phages and their bacterial hosts can be easily propagated in controlled laboratory environments where short generation times favour rapid evolutionary rates. Moreover, lytic phages in particular (i.e. those that must kill their host cell to reproduce), have simple life histories and relatively small, well-understood genomes allowing reliable mapping of genotype to phenotype and the formulation of a detailed knowledge of the pathways to adaptation. Importantly, phages are themselves of ecological and economic importance, for example, in structuring natural bacterial communities [2] and for their undesirable effects on bioprocessing of food and waste products by bacteria [3]. As a result of these factors, phages have emerged as a key model system for the study of viral evolution and in particular viral host-adaptation [1]. To date a wide range of phages and their associated bacterial hosts have been employed in experimental evolution studies encompassing a broad sweep of phylogenetic diversity and different forms of genomic organisation [1,4]. In this article we summarise recent advances in our understanding of viral host-adaptation arising from experimental evolution of phages. We highlight the differences in the pattern and process of phage evolution depending upon whether adaptation is against a fixed or a coevolving host, and outline the genetic and ecological factors that shape the evolution of phage host range.

Experimental evolution

As with all viruses, phage replication depends on intimate interactions with a number of host cell components, from receptors that mediate entry, to transcription and translation

49 machinery that produce virus particles, to cell wall components that are disrupted to lyse
50 the host. Evolution can rapidly tune the kinetics of these reactions, and the huge
51 population sizes that phage can reach, combined with the frequency of mutation and
52 their generally compact genomes, ensures that there is abundant variation on which
53 selection can efficiently act [5]. When repeatedly exposed to a given host genotype,
54 phages tend to evolve higher growth rates [6-11] and/or an increased phage-imposed
55 reduction in host growth [8,12] when compared with their ancestors (Figure 1). Many
56 genes, including those involved in capsid structure [11,13], tail fibres [14] and viral
57 replication [13], have been identified as undergoing changes during adaptation to a host,
58 although many studies have been conducted with confounding factors (such as
59 increased temperature, or growth in a chemostat) making it difficult to identify the genetic
60 targets specific to host adaptation *per se*. A principal feature of phage host-adaptation is
61 that evolution can be highly parallel, with fixation of the same mutations occurring in
62 independently evolving populations of phage. Indeed, 'replaying the tape' of evolution
63 often has repeatable consequences [13], and 'rewinding the tape' by reversing the
64 selective pressures (i.e. adapting back to the host of the ancestral phage) sees the
65 appearance of mutations reverting the phage genome towards the ancestral state
66 [6,15,16]. Where adaptation does not necessarily require many amino acid changes
67 [7,17] and there are limits to the number of beneficial mutations available due to the
68 small size and high levels of pleiotropy in a virus genome [18], parallel evolution might
69 be expected [19,20]. (Pleiotropy refers to the case where a single gene affects multiple
70 functions.) Interestingly, parallel 'silent' mutations (those that do not affect the amino
71 acid sequence of translated virus products) have been frequently observed, indicating
72 that selection can be strong on features other than translated gene products, such as
73 nucleic acid secondary structure or translation efficiency [6,7,11].

The implication of parallel adaptation is that the mutations involved have consistent beneficial effects on fitness, as phages optimise their interactions with host processes in the context of their own genetics and ecology. However, these fitness effects can be highly specific, with different adaptive paths open to different genotypes. Individual mutations may not act independently, but rather be dependant on the pre-existing genetic background (i.e. epistasis) [7,10]. When three related species of phage were adapted to a host, parallel evolution within each species was high, but the different species did not acquire similar mutations at homologous sites as they adapted [21]. Epistasis, coupled with an incomplete understanding of the precise structures and processes involved in the phage lifecycle, mean that whilst qualitative phenotypic trends occurring as a phage adapts to its host tend to be predictable, anticipating the exact effects of selection in shaping phenotype or genotype has proved somewhat difficult. For example, adaptation to altered host receptor lipopolysaccharide resulted in changes to the ϕ X174 virus capsid as predicted, but these changes were located internally, not in the receptor binding site, and seemed to play a more general role in enhancing capsid stability than receptor specificity [11]. Likewise, selection under conditions favouring an altered lysis time yielded phages with phenotypes qualitatively in line with predictions but significantly divergent from the values predicted by a theoretical model [22,23]. Nevertheless, the fact that phage fitness tends to plateau under consistent selection [6,7,10] points to a dynamic in which evolution on a fixed host is selecting for an optimal phenotype in the context of the experimental conditions, towards which the population converges.

Evolving optimum fitness

Several obstacles may prevent viruses from attaining this optimum. Some traits may

99 require many simultaneous mutations to evolve, or are constrained by the physical
100 properties of the genes and proteins involved, and thus are unlikely to appear in small
101 populations or over short periods of time. Population bottlenecks, caused by events such
102 as transmission, dispersal, and population dynamics often occur during virus evolution
103 [24-26] and have the potential to shape the course of evolution by preventing the
104 appearance and spread of beneficial mutations and facilitating fixation of deleterious
105 ones [15,27]. Similarly, neutral and slightly deleterious mutations can hitchhike to fixation,
106 if they fortuitously occur alongside a beneficial mutation in an expanding clone. The high
107 mutation rate and small genome size, particularly of RNA viruses, can potentially
108 overcome these problems by replenishing population diversity, allowing even a clonal
109 virus to overcome the genetic restrictions of a bottleneck and efficiently explore the local
110 fitness landscape [28]. A balance must be struck, however, because an elevated rate of
111 mutation can inhibit optimal fitness by consistently mutating viruses away from a fitness
112 peak [29], which in combination with frequent bottlenecks can lead to lethal mutagenesis
113 and extinction [29].

114 Viruses within a population can also compete with one another. Clonal interference,
115 whereby the rate of adaptation is inhibited by competition between different beneficial
116 mutations, can be a major impediment to adaptation [30] although at high mutation rates
117 and population size its effects can be somewhat alleviated by mutational supply [31].
118 Where supply of hosts is limited, multiplicity of infection (MOI) is high, and virus clones
119 can co-infect the same host. Co-infection can allow for recombination, potentially
120 increasing the rate of adaptation (although this effect can be small if co-infecting phages
121 are closely related [31]. More importantly perhaps, co-infecting virus clones must
122 compete for host resources. Under such conditions, there may not be a universal
123 optimum genotype. In some cases, virus clones can parasitize the genes of co-infecting

viruses, resulting in a 'prisoner's dilemma' where the fitness of a particular genotype is dependent on the frequency of competitors [32]. Phages can even evolve the means to detect co-infection, and modulate their infection strategy accordingly. Phage $\phi 2$ evolved under high MOI killed *Pseudomonas fluorescens* host cells more rapidly than those evolved under low MOI, but only when assay MOI was high [33]. Such adaptive phenotypic plasticity raises the possibility that life-history plasticity is a trait amenable to selection in viruses. The tensions resulting from these competing interests mean that the dynamics of these relationships are rapid and unpredictable, and virus evolution can continue even in long-term cultures [13].

Experimental coevolution

In contrast to the optimisation of viral fitness achievable during experimental evolution, scenarios where bacteria too are allowed to evolve - i.e. experimental coevolution [34] - may preclude net gains in viral fitness due to the potential for continual reciprocal adaptation and counter-adaptation inherent to antagonistic coevolution. Indeed, empirical studies, across a range of bacteria-phage systems, reveal that rapid, persistent antagonistic coevolution is a common outcome of co-propagation of bacteria and phage [35-39] (challenging the view that bacteria-phage coevolution is universally constrained [40]). These studies suggest that bacteria-phage coevolution typically takes the form of an arms-race of repeated cycles of evolution of bacterial-resistance followed by evolution of phage to infect these resistant bacteria, which can last for several hundreds of bacterial generations [35,37,41]. This process is driven by predominantly directional selection favouring the evolution of broader virus host-range and, concomitantly, broader bacterial resistance range through time, but no corresponding increase in viral growth rate on the ancestral host [8] (Figure 1). As such, for a given population, phages from

later in the experiment can typically infect bacterial genotypes from the past, even though these bacteria will tend to be resistant to their own contemporaneous phages [37,39,42].

Recurrent selective sweeps of new infectivity mutations leads to the accumulation of multiple infectivity mutations per phage genome, and a tendency for host-range and the number of substitutions to be positively correlated [37,42]. Indeed, this stepwise accretion of mutations appears to be crucial to effective host-range expansion [12,17]. Continual reciprocal selection for innovation tends to accelerate viral evolution in coevolving populations compared to viral populations adapting to a fixed host environment [14,38], in some cases as much as doubling the nonsynonymous substitution rate [14]. Moreover, coevolution leads to greater between-population divergence of viral genomes than does selection against a fixed host [14]. Thus, while common loci are targeted by selection across replicate coevolving populations, the specific sites of mutations and/or the combinations of co-occurring mutations vary, giving rise to phenotypic divergence in viral host-range between populations [14] (Figure 1). In some phages, host-range mutations have been observed in genes known to encode infectivity determinants such as proteins for host-binding (e.g. tail fibre protein in phage $\phi 2$ [14,42], host recognition protein J in phage lamda [17], whereas, in other phages, host-range mutations have been observed in genes of unknown function, suggesting novel mechanisms of bacteria-phage interaction [37].

Host-range expansion & host-shifts

The expansion or shift of viral host range is one of the most pertinent features of viral evolution. Many recent examples of emerging human viral diseases, such as HIV, bird/swine flu and SARS are the result of expanding host range [1]. However, co-culture

174 studies have demonstrated that host range shifts are generally highly constrained by
175 both genetic and ecological factors.

176 While coevolution leads to the stepwise build-up of broad host-range, larger shifts onto
177 more distantly related hosts presents a greater challenge for viruses. Attempts to
178 experimentally evolve phages to infect novel hosts often fail, even when the new host is
179 of the same species [12,43]. Where larger numbers of mutations are required to infect a
180 novel host, especially where those mutations act synergistically, the likelihood of
181 evolving infectivity in one step is greatly reduced. For example, when exposed to
182 resistant *P. fluorescens* strains (derived from coevolution with phage $\phi 2$), the ancestral
183 phage genotype could evolve to infect hosts in one step, but only when relatively few
184 mutations were required [12]. Interestingly, sequencing of host range mutants has
185 shown that host range evolution is almost always associated with mutations in specific
186 host attachment proteins (tail-fibre proteins in $\phi 2$ [12] and T7 [44], and attachment
187 protein P3 in $\phi 6$ [43,45], suggesting a relatively limited array of virus genes on which
188 selection can act. Access to the native host, either through migration [46] or in a mixed
189 host environment [47,48], can greatly increase the potential for host range shifts.
190 Susceptible hosts support a 'source' population for host range mutants able to infect
191 novel hosts with poor efficiency ('sink' populations). Source populations provide the
192 means to maintain population size and thus mutation supply, increasing the opportunity
193 for beneficial mutations to arise.

194 Where host range mutations do arise they often result in antagonistic pleiotropic effects,
195 reducing fitness in the native host [8,43,45], or even causing a loss of infectivity [9].
196 Evolution of host range is therefore likely to be subject to a trade-off in the benefits of
197 host range expansion vs. specialisation. Phage selection experiments have
198 demonstrated the role of several key ecological variables that shape the outcome of host

range evolution. Firstly, availability of susceptible hosts during adaptation to a novel host can hinder, as well as facilitate, the evolution of range shifts [48]. When evolved in varying ratios of susceptible:novel hosts, phage $\phi 2$ only evolved to infect a novel host when the frequency of susceptible hosts was between 0.1 – 1% relative to the novel host [48]. Phage genotypes able to grow on both the native and novel hosts were shown to have a significantly lower growth rate on each host population compared with host specialists. Thus, in the presence of greater frequencies of susceptible hosts, even where novel hosts remain in the majority, the pleiotropic cost of expanded host range results in selection against generalists. Furthermore, where host range expansions did occur, the vast majority of host range mutants evolved to specialise on the novel host alone, resulting in a corresponding increase in fitness on the new host. A similar pattern of adaptation was observed in the $\phi 6$ phage of *Pseudomonas syringae* [9], suggesting that even where host range shifts are favoured, antagonistic pleiotropy will favour divergent evolution of host specialists rather than the stable existence of generalists.

Secondly, host range shifts are also dependent on the intrinsic quality of the novel host resource. Under optimal foraging theory, where there is a disparity in the profitability of different hosts, specialism on the more profitable host will be favoured. For example, populations of a generalist T7 phage able to infect two *E. coli* host strains, one strain of which was less profitable than the other, evolved to infect only the more profitable host [44]. Crucially, selection for optimal foraging is not dependant on a cost of generalism. Indeed, in this study avoidance of the poor quality host was associated with a drop in infectivity on the preferred host [44].

Thirdly, Bono et al. have shown that intraspecific competition for resources can be a major driving force behind the evolution of generalism. Increasing the MOI of $\phi 6$ phages in *P. syringae* populations lead to an increase in the probability of generalist emergence

[49]. Furthermore, the authors find that the impact of intraspecific competition was positively associated with the quality of the novel host, driving rapid host range expansions where competition is strong and host quality is relatively equal, and slow rates of host range evolution where novel hosts are considerably less profitable than the native host [49].

Concluding remarks

Experimental studies of phage evolution have yielded substantial advances in our understanding of viral host-adaptation. Several clear trends emerge: First, viral adaptation to a given host genotype often results in parallel evolution and the convergence of independent lineages upon a shared adaptive solution via the same genetic targets, although this parallelism does not appear to be conserved when comparing evolution of different phage species. Second, reciprocal evolution of the bacterial host tends to accelerate phage evolution through continual selection for increased infectivity and leads to greater between-population divergence. Third, phage host-range expansion occurs during coevolution by stepwise evolution involving multiple mutations, and large host-shifts appear to be constrained by mutational accessibility, pleiotropic costs and ecological factors.

It is interesting to consider the extent to which these evolutionary patterns translate to viruses of eukaryotes. First, there are clear and important biological differences between prokaryotic and eukaryotic hosts, for example, multicellular eukaryotes tend to live in much smaller populations than bacteria, reproduce more slowly and only a small fraction of host cells are germline. These differences are likely to generate contrasting selection and demand different adaptive solutions. Nevertheless, several aspects of studies on viruses and their eukaryotic hosts appear consistent with the general patterns outlined

above. For example, as with phages, adaptation to a particular host favoured parallel adaptations in experimental evolution of influenza virus A [50] and tobacco etch potyvirus (TEV) [51]. Antagonistic pleiotropy and multiple steps to host range expansion have been recorded in experimental TEV evolution [51] and in the history of feline panleukopenia virus evolution [52]. Although rapid coevolution between viruses and their eukaryotic hosts is constrained at the organismal level by the relatively slow eukaryote rate of replication (and hence coevolution), coevolutionary dynamics are apparent in the interaction between virus serotypes and the adaptive immunity and immune memory of vertebrates. As with bacteria-phage coevolution, this can lead to rapid evolution and between-population divergence, as has been observed in the case of HIV-antibody coevolution within a patient [53] and the ‘antigenic drift’ of foot-and-mouth disease virus [54] and influenza virus [55] between host populations.

Numerous questions remain for future research. Existing studies have largely focused on phage adaptation at the interface of virus-host attachment, yet the recent discovery of CRISPR-based immunity suggests huge potential for dynamic intracellular responses by bacteria to act as a driver of phage evolution (e.g. see [56,57]). Very few studies have addressed the extent to which the evolutionary characteristics of phages are phylogenetically conserved (although see [21]) suggesting exciting potential for ‘comparative’ experimental evolution. In particular, it is unclear why some bacteria-phage interactions undergo extensive, prolonged arms race coevolution whereas others apparently do not (e.g., [58,59]). While host-adaptation by lytic phages has been well studied, temperate phages have not been extensively studied in an experimental evolution context (although see [60]). Finally, natural phage communities are highly diverse, thus it is important to understand how the evolution of phages in isolation scales up when embedded in more complex, species rich phage communities.

274

275 **Acknowledgements**

276 Our work is supported by grants from the European Research Council (COEVOCON
277 311490) and the Natural Environment Research Council (NE/H005080/1) to M.A.B.

278

279 **References**

- 280 1. Dennehy JJ: **Bacteriophages as model organisms for virus emergence**
281 **research.** *Trends Microbiol* (2009) **17**(10):450-457.
282
- 283 2. Weinbauer MG: **Ecology of prokaryotic viruses.** *FEMS Microbiol Rev* (2004)
284 **28**(2):127-181.
285
- 286 3. Los M: **Minimization and prevention of phage infections in bioprocesses.**
287 *Methods Mol Biol* (2012) **834**(305-315).
288
- 289 4. Dennehy JJ: **What can phages tell us about host-pathogen coevolution?** *Int*
290 *J Evol Biol* (2012) **2012**(396165).
291
- 292 5. Elena S, F. , Sanjuan R: **Virus evolution: Insights from an experimental**
293 **approach.** *Annual Review of Ecology, Evolution, and Systematics* (2007) **38**(
294
295 An excellent review of viral evolution with a focus on mutation rates and the evolution of
296 robustness, and the key differences between RNA and DNA-based viral
297 genomes
298
- 299 6. Bull J, Badgett M, Wichman H, Huelsenbeck J, Hillis D, Gulati A, Ho C, Molineux
300 I: **Exceptional convergent evolution in a virus.** *Genetics* (1997) **147**(4):1497-
301 1507.
302
- 303 7. Wichman H, Badgett M, Scott L, Boulianne C, Bull J: **Different trajectories of**
304 **parallel evolution during viral adaptation.** *Science* (1999) **285**(5426):422-424.
305
- 306 The classic study demonstrating the propensity for parallel molecular evolution in
307 phages
308
- 309 8. Poullain V, Gandon S, Brockhurst M, Buckling A, Hochberg M: **The evolution of**
310 **specificity in evolving and coevolving antagonistic interactions between a**
311 **bacteria and its phage.** *Evolution; international journal of organic evolution*
312 (2008) **62**(1):1-11.
313
- 314 9. Duffy S, Burch CL, Turner PE: **Evolution of host specificity drives**
315 **reproductive isolation among rna viruses.** *Evolution* (2007) **61**(11):2614-2622.

- 316
317 10. Nguyen A, Molineux I, Springman R, Bull J: **Multiple genetic pathways to**
318 **similar fitness limits during viral adaptation to a new host.** *Evolution;*
319 *international journal of organic evolution* (2012) **66**(2):363-374.
320
321 11. Pepin K, Lambeth K, Hanley K: **Asymmetric competitive suppression**
322 **between strains of dengue virus.** *BMC Microbiol* (2008) **8**(28).
323
324 12. Hall AR, Scanlan PD, Buckling A: **Bacteria-phage coevolution and the**
325 **emergence of generalist pathogens.** *Am Nat* (2011) **177**(1):44-53.
326
327 13. Wichman HA: **Adaptive molecular evolution for 13,000 phage generations: A**
328 **possible arms race.** *Genetics* (2005) **170**(
329
330 14. Paterson S, Vogwill T, Buckling A, Benmayor R, Spiers AJ, Thomson NR, Quail
331 M, Smith F, Walker D, Libberton B, Fenton A *et al*: **Antagonistic coevolution**
332 **accelerates molecular evolution.** *Nature* (2010) **464**(7286):275-278.
333
334 Demonstrates that host evolution accelerates phage evolution, also leading to greater
335 population divergence and higher levels of within population diversity
336
337 15. Turner P, McBride R, Duffy S, Montville R, Wang L-S, Yang Y, Lee S, Kim J:
338 **Evolutionary genomics of host-use in bifurcating demes of rna virus phi-6.**
339 *BMC Evol Biol* (2012) **12**(153).
340
341 16. Crill W, Wichman H, Bull J: **Evolutionary reversals during viral adaptation to**
342 **alternating hosts.** *Genetics* (2000) **154**(1):27-37.
343
344 17. Meyer J, Dobias D, Weitz J, Barrick J, Quick R, Lenski R: **Repeatability and**
345 **contingency in the evolution of a key innovation in phage lambda.** *Science*
346 *(New York, NY)* (2012) **335**(6067):428-432.
347
348 18. Belshaw R, Gardner A, Rambaut A, Pybus O: **Pacing a small cage: Mutation**
349 **and rna viruses.** *Trends Ecol Evol* (2008) **23**(4):188-193.
350
351 19. Orr H: **The probability of parallel evolution.** *Evolution; international journal of*
352 *organic evolution* (2005) **59**(1):216-220.
353
354 20. Wood T, Burke J, Rieseberg L: **Parallel genotypic adaptation: When evolution**
355 **repeats itself.** *Genetica* (2005) **123**(1-2):157-170.
356
357 21. Bollback J, Huelsenbeck J: **Parallel genetic evolution within and between**
358 **bacteriophage species of varying degrees of divergence.** *Genetics* (2009)
359 **181**(1):225-234.
360
361 One of the few comparative studies of related phages suggests that patterns of parallel
362 evolution are restricted and do not extend between virus species
363
364 22. Heineman R, Bull J: **Testing optimality with experimental evolution: Lysis**
365 **time in a bacteriophage.** *Evolution; international journal of organic evolution*
366 (2007) **61**(7):1695-1709.

23. Chanturanupong L, Heineman R: **A common, non-optimal phenotypic endpoint in experimental adaptations of bacteriophage lysis time.** *BMC Evol Biol* (2012) **12**(37).
24. Forrester NL, Guerbois M, Seymour RL, Spratt H, Weaver SC, Vignuzzi M: **Vector-borne transmission imposes a severe bottleneck on an rna virus population.** *PLoS Pathogens* (2012) **8**(
25. Domingo E, Escarmis C, Lazaro E, Manrubia S: **Quasispecies dynamics and rna virus extinction.** *Virus Res* (2005) **107**(2):129-139.
26. Li H, Roossinck M: **Genetic bottlenecks reduce population variation in an experimental rna virus population.** *J Virol* (2004) **78**(19):10582-10587.
27. Patwa Z, Wahl L: **The impact of host-cell dynamics on the fixation probability for lytic viruses.** *J Theor Biol* (2009) **259**(4):799-810.
28. Burch C, Chao L: **Evolution by small steps and rugged landscapes in the rna virus phi6.** *Genetics* (1999) **151**(3):921-927.
29. Santiago FE, Rafael Sn: **Virus evolution: Insights from an experimental approach.** *Annual Review of Ecology, Evolution, and Systematics* (2007) **38**(
30. Miralles R: **Clonal interference and the evolution of rna viruses.** *Science* (1999) **285**(
31. Bollback JP, Huelsenbeck JP: **Clonal interference is alleviated by high mutation rates in large populations.** *Mol Biol Evol* (2007) **24**(
32. Turner PE, Chao L: **Prisoner's dilemma in an rna virus.** *Nature* (1999) **398**(6726):441-443.
33. Leggett H, Benmayor R, Hodgson D, Buckling A: **Experimental evolution of adaptive phenotypic plasticity in a parasite.** *Current biology : CB* (2013) **23**(2):139-142.
- Demonstrates the evolution of plasticity in a phage whereby lysis time varies according to coinfection status
34. Brockhurst MA, Koskella B: **Experimental coevolution of species interactions.** *Trends Ecol Evol* (2013) **28**(6):367-375.
35. Buckling A, Rainey PB: **Antagonistic coevolution between a bacterium and a bacteriophage.** *Proceedings of the Royal Society of London Series B-Biological Sciences* (2002) **269**(1494):931-936.
36. Mizoguchi K, Morita M, Fischer CR, Yoichi M, Tanji Y, Unno H: **Coevolution of bacteriophage pp01 and escherichia coli o157:H7 in continuous culture.** *Appl Environ Microb* (2003) **69**(1):170-176.

37. Marston MF, Pierciey FJ, Shepard A, Gearin G, Qi J, Yandava C, Schuster SC, Henn MR, Martiny JBH: **Rapid diversification of coevolving marine synechococcus and a virus.** *P Natl Acad Sci USA* (2012) **109**(12):4544-4549.
38. Kashiwagi A, Yomo T: **Ongoing phenotypic and genomic changes in experimental coevolution of rna bacteriophage qbeta and escherichia coli.** *Plos Genet* (2011) **7**(8):e1002188.
39. Meyer JR, Flores C, Weitz JS, Lenski RE: **Key innovation in a virus catalyzes a coevolutionary arms-race.** *13:ALife* 532-533.
40. Lenski RE, Levin BR: **Constraints on the coevolution of bacteria and virulent phage - a model, some experiments, and predictions for natural communities.** *Am Nat* (1985) **125**(4):585-602.
41. Hall AR, Scanlan PD, Morgan AD, Buckling A: **Host-parasite coevolutionary arms races give way to fluctuating selection.** *Ecol Lett* (2011) **14**:635-642.
42. Scanlan PD, Hall AR, Lopez-Pascua LDC, Buckling A: **Genetic basis of infectivity evolution in a bacteriophage.** *Mol Ecol* (2011) **20**(5):981-989.
43. Duffy S, Turner PE, Burch CL: **Pleiotropic costs of niche expansion in the rna bacteriophage phi 6.** *Genetics* (2006) **172**(2):751-757.
44. Heineman RH, Springman R, Bull JJ: **Optimal foraging by bacteriophages through host avoidance.** *Am Nat* (2008) **171**(4):E149-E157.
- Demonstrates that phages evolve to avoid low quality hosts and even do so despite incurring a cost to their reproduction in high quality hosts
45. Ferris M, Joyce P, Burch C: **High frequency of mutations that expand the host range of an rna virus.** *Genetics* (2007) **176**(2):1013-1022.
46. Ching J, Musheyev S, Chowdhury D, Kim J, Choi Y, Dennehy J: **Migration enhances adaptation in bacteriophage populations evolving in ecological sinks.** *Evolution; international journal of organic evolution* (2013) **67**(1):10-17.
47. Dennehy JJ, Friedenber NA, Holt RD, Turner PE: **Viral ecology and the maintenance of novel host use.** *Am Nat* (2006) **167**(3):429-439.
48. Benmayor R, Hodgson DJ, Perron GG, Buckling A: **Host mixing and disease emergence.** *Curr Biol* (2009) **19**(9):764-767.
- Demonstrates that that the ecological conditions promoting phage host-shifts are highly restrictive
49. Bono LM, Gensel CL, Pfennig DW, Burch CL: **Competition and the origins of novelty: Experimental evolution of niche-width expansion in a virus.** *Biology Letters* (2013) **9**(1).
50. Keleta L, Ibricevic A, Bovin NV, Brody SL, Brown EG: **Experimental evolution**

of human influenza virus h3 hemagglutinin in the mouse lung identifies adaptive regions in ha1 and ha2. *J Virol* (2008) 82(23):11599-11608.

51. Bedhomme S, Lafforgue G, Elena SF: **Multihost experimental evolution of a plant rna virus reveals local adaptation and host-specific mutations. *Mol Biol Evol* (2012) 29(5):1481-1492.**
52. Shackelton LA, Parrish CR, Truyen U, Holmes EC: **High rate of viral evolution associated with the emergence of carnivore parvovirus. *Proc Natl Acad Sci U S A* (2005) 102(2):379-384.**
53. Liao H-X, Lynch R, Zhou T, Gao F, Alam SM, Boyd SD, Fire AZ, Roskin KM, Schramm CA, Zhang Z, Zhu J *et al*: **Co-evolution of a broadly neutralizing hiv-1 antibody and founder virus. *Nature* (2013) 496(7446):469-+.**
54. Borrego B, Novella IS, Giralt E, Andreu D, Domingo E: **Distinct repertoire of antigenic variants of foot-and-mouth-disease virus in the presence or absence of immune selection. *J Virol* (1993) 67(10):6071-6079.**
55. Carrat F, Flahault A: **Influenza vaccine: The challenge of antigenic drift. *Vaccine* (2007) 25(39-40):6852-6862.**
56. Levin BR, Moineau S, Bushman M, Barrangou R: **The population and evolutionary dynamics of phage and bacteria with crispr-mediated immunity. *Plos Genet* (2013) 9(3):e1003312.**
57. Sun CL, Barrangou R, Thomas BC, Horvath P, Fremaux C, Banfield JF: **Phage mutations in response to crispr diversification in a bacterial population. *Environ Microbiol* (2013) 15(2):463-470.**
58. Brockhurst MA, Buckling A, Rainey PB: **Spatial heterogeneity and the stability of host-parasite coexistence. *J Evolution Biol* (2006) 19(2):374-379.**
59. Lythgoe KA, Chao L: **Mechanisms of coexistence of a bacteria and a bacteriophage in a spatially homogeneous environment. *Ecol Lett* (2003) 6(4):326-334.**
60. Refardt D, Rainey PB: **Tuning a genetic switch: Experimental evolution and natural variation of prophage induction. *Evolution* (2010) 64(4):1086-1097.**

An elegant study of optimal life history evolution in a temperate phage

Figure Legend

Figure 1. Contrasting trajectories of viral adaptation in terms of growth rate and host-range phenotypes for viruses evolving against a fixed host genotype (E - i.e. 'evolving')

516 or a coevolving host population (C – i.e. ‘coevolving’). Arrows show the trajectory of
517 evolution from the ancestral phenotype (A – i.e. ‘ancestral’) and while trajectories are
518 parallel for evolving viral lineages, they are divergent for coevolving viral lineages
519 (compare ‘E’ and ‘C’ trajectories). Adapted from data presented in ref. 8.

